

---

Travaux Dirigés de Physique des Particules

---

**QED :  $e^+e^-$  Bhabha scattering**  
Feynman Diagrams, Feynman Rules, cross-section calculation

---

**Reminder : QED Feynman Rules**

These rules, originated from Quantum Field Theory, are applied on Feynman diagrams in order to evaluate the transition matrix element. For the external lines of a diagram :

- Each incoming spin  $\frac{1}{2}$  particle is associated to :  $u(p, s)$
- Each outgoing spin  $\frac{1}{2}$  particle is associated to :  $\bar{u}(p, s)$
- Each incoming spin  $\frac{1}{2}$  antiparticle is associated to :  $\bar{v}(p, s)$
- Each outgoing spin  $\frac{1}{2}$  antiparticle is associated to :  $v(p, s)$
- Each incoming photon is associated to :  $\varepsilon_\mu(p, \lambda)$
- Each outgoing photon is associated to :  $\varepsilon_\mu^*(p, \lambda)$

For the internal lines of a diagram :

- Photon propagator :  $\frac{-ig^{\mu\nu}}{q^2}$
- Fermion propagator :  $\frac{i\gamma^\mu p_\mu + m}{q^2 - m^2}$
- Massive boson propagator :  $\frac{i}{q^2 - m^2}$

For a vertex between a photon and to charge  $e$  fermions :  $-ie\gamma^\mu$


In general there is more than one Feynman diagram contributing to a process and sometimes identical particles appear in the initial or final states.

- Write all topological inequivalent diagrams at any order in perturbation theory.
  - The overall sign of a given diagram is not observable, but diagrams that differ in the exchange of two identical fermions in the initial or final state, or a fermion-antifermion in the initial or final state, respectively, should come with opposite signs on account of Fermi statistics.
-

**Problem : Bhabha scattering :**  $e^+e^- \longrightarrow e^+e^-$

**1 /** Draw the Feynman diagrams contributing to this scattering. In this problem, we restraint our study to the dominant processes at low energy.

**2 /** The kinetic variables are defined in the following figure. Give the Mandelstam variables  $s = (k + p)^2$ ,  $t = (k - k')^2$  and  $u = (k - p')^2$  in the ultra relativist limit.



**3 /** Use Feynman rules to calculate the transition matrix element  $T$ , and its conjugate  $T^*$

**4 /** Show that  $|T|^2$  is the sum of three contributions : scattering ( $T_t$ ), annihilation ( $T_s$ ) and interference.

**5 /** Show that the contribution of the annihilation term averaged on the initial polarizations and summed on the final polarizations is:

$$\overline{|T_s|^2} = \frac{e^4}{4q^4} \text{Tr}((\not{k} - m_e)\gamma_\mu(\not{p} + m_e)\gamma_\nu) \text{Tr}((\not{p}' + m_e)\gamma^\mu(\not{k}' - m_e)\gamma^\nu)$$

**6 /** Calculate  $\overline{|T_s|^2}$  in the ultra-relativist limit.

Reminder : some properties of the  $\gamma$  matrices :

$$\text{Tr}(\gamma^\mu\gamma^\nu\gamma^\alpha\gamma^\beta) = 4(g^{\mu\nu}g^{\alpha\beta} - g^{\mu\alpha}g^{\nu\beta} + g^{\mu\beta}g^{\nu\alpha})$$

$$\gamma_\alpha\gamma_\mu\gamma_\nu\gamma^\alpha = 4g_{\mu\nu}$$

$$\gamma_\alpha\gamma_\mu\gamma_\nu\gamma_\rho\gamma^\alpha = -2\gamma_\rho\gamma_\nu\gamma_\mu$$

**7 /** Repeat step 5 and 6 to calculate the contributions of the scattering and the interference terms.

**8 /** Show that the Bhabha differential cross-section is :

$$\frac{d\sigma}{d\cos\theta} = \frac{\pi\alpha^2}{s} \left( \frac{s^2 + u^2}{t^2} + \frac{2u^2}{ts} + \frac{u^2 + t^2}{s^2} \right)$$

**9 /** Let  $\theta$  be the scattering angle in the center of mass frame. Show that in this frame :

$$t = -\frac{s}{2}(1 - \cos\theta) \quad \text{and} \quad u = -\frac{s}{2}(1 + \cos\theta)$$

Give the Bhabha cross section in the center of mass frame.

**10 /** How to calculate the Møller scattering cross-section ( $e^-e^- \longrightarrow e^-e^-$ ) from the previous results ? Show that

$$\frac{d\sigma}{d\cos\theta} = \frac{\pi\alpha^2}{s} \left( \frac{s^2 + u^2}{t^2} + \frac{2s^2}{tu} + \frac{s^2 + t^2}{u^2} \right)$$