

Chapter 6

∞

Electromagnetic interactions

Outline/Plan

1. Cross-section
2. Matrix element
3. Quantum Electrodynamics
(QED) Feynman rules

1. Section efficace
2. Élément de matrice
3. Règles de Feynman pour
l'électro-dynamique quantique

1. Cross-section

Consider the general process for two-particle scattering :

$$a+b \rightarrow 1+2+\dots+n$$

- States normalization : in NRQM usually one takes 1 particle per unit volume. For KG solutions :

$$\phi_a = N_a e^{-ip_a x} \Rightarrow \rho_a = i \left[\phi_a^* (\partial_t \phi_a) - (\partial_t \phi_a^*) \phi_a \right] = 2N_a^2 E_a$$

- Usual prescription (covariant normalization) :

2E particles per unit volume

$$\int_V d^3x \rho_a = 2E_a \Rightarrow N_a = V^{-1/2}$$

1. Cross-section

The general expression of the cross-section reads :

$$d\sigma = \frac{1}{2E_a 2E_b |\vec{v}_a - \vec{v}_b|} \int (2\pi)^4 \delta^4(p_f - p_i) \overline{|T_{fi}|^2} \prod_{k=1}^{k=n} \frac{d^3 p_k}{(2\pi)^3 2E_k}$$

- $2E_a 2E_b |\vec{v}_a - \vec{v}_b| / V^2$ is the flux factor (number of incident particles per unit area reaching the target per unit time \times number of target particles) with normalization at $2E$ particles per unit volume.
- $\prod_{k=1}^{k=n} \frac{V d^3 p_k}{(2\pi)^3 2E_k}$ is the final state available phase space (for one particle in a volume V with a momentum in $d^3 p$ one has the usual expression : $V d^3 p / (2\pi)^3$)

1. Cross-section

- $(2\pi)^4 \delta^4(p_f - p_i) \overline{|T_{fi}|^2}$ is the transition rate per volume unit. It is the part containing the physics of the interaction processes. Each diagram's line is associated with a normalization factor N_i ($i=a,b,1,\dots,n$) which induces an overall factor :

$$\left[\underbrace{(V^{-1/2})^2}_{a,b} \times \underbrace{(V^{-1/2})^n}_{1\dots n} \right]^2 = V^{-(n+2)}$$

balancing the contributions of the flux and the phase space => the cross-section is independent of the normalization volume!

1. Cross-section

Application : Rutherford scattering formula (2+2 process => n=2).

- Flux factor : $2E_a 2E_b |\vec{v}_a - \vec{v}_b| = 4 \left[(p_a \cdot p_b)^2 - m_a^2 m_b^2 \right] = 4 |\vec{p}_a^*| \sqrt{s}$
where the Mandelstam variable s has been used $s = (p_a + p_b)^2$

- 2-bodies phase space :


$$\begin{aligned}
 dQ_2 &= \int_4 (2\pi)^4 \delta^4(p_f - p_i) \frac{d^3 p_1}{(2\pi)^3 2E_1} \frac{d^3 p_2}{(2\pi)^3 2E_2} \\
 &= \frac{1}{16\pi^2} \frac{|\vec{p}_1|^3 d\Omega_1}{|\vec{p}_1|^2 E_T - \vec{p}_1 \cdot \vec{p}_T E_1} \quad \text{with } (p_T) = (p_a + p_b) \\
 &= \frac{1}{16\pi^2} \frac{|\vec{p}_1^*| d\Omega_1}{\sqrt{s}}
 \end{aligned}$$

1. Cross-section

Application : Rutherford scattering formula (2+2 process => n=2).

- Transition rate :

$$|T_{fi}|^2 = \left(\frac{e^2}{q^2} \right)^2 (p_a + p_b) \cdot (p_1 + p_2)$$



- Mandelstam variables :

$$s = (p_a + p_1)^2 \quad t = (p_a - p_b)^2 \quad u = (p_a - p_2)^2$$

$$|T_{fi}|^2 = \left(\frac{4\pi\alpha}{t} \right)^2 (s - u)^2 \quad \text{and} \quad \frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2 s} \frac{|\vec{p}_1^*|}{|\vec{p}_a^*|} |T_{fi}|^2$$

- In the LAB frame (target at rest)

1. Cross-section

Application : Rutherford scattering formula (2+2 process => n=2).

- “Classical” case : α is a light particle (negligible m), 1 at rest initially => use of the LAB frame

$$p_a \simeq \left(|\vec{k}|, \vec{k} \right) \quad p_1 \simeq \left(M, \vec{0} \right)$$

$$t \simeq -4 |\vec{k}|^2 \sin^2(\theta/2) \quad (s-u)^2 = 16 |\vec{k}|^2 M^2$$

$$4 \left[(p_a \cdot p_1)^2 - m^2 M^2 \right] \simeq 4 |\vec{k}|^2 M^2$$


- With the fine structure constant $\alpha = e^2 / 4\pi$ one gets the standard Rutherford formula :

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4 |\vec{k}|^2 \sin^4(\theta/2)}$$

1. Cross-section

Question : how one computes the transition rate

$$|T_{fi}|^2 = \left(\frac{e^2}{q^2} \right)^2 (p_a + p_1) \cdot (p_b + p_2)$$



?


2- Matrix element

- Classical theory : transition amplitude (covariant expression) from initial (i) to final (f) state under the action of a perturbation potential V .

$$T_{fi} = -i \int d^4x \phi_f^*(x) V(x) \phi_i(x)$$

- 1st order K.G. potential : $V = -ie(\partial_\mu A^\mu + A^\mu \partial_\mu)$
- Integrating by parts introduces the transition current :


$$T_{fi} = -i \int d^4x A^\mu \underbrace{(-ie) [\phi_f^* \partial_\mu \phi_i - \partial_\mu \phi_f^* \phi_i]}_{J_\mu^{(fi)}}$$



2. Matrix element


- (i) and (f) are taken as free particle solutions : $\phi_{i,f}(x) = N_{i,f} e^{-ip_{i,f}x}$
- Inserting the free particle solutions leads to

$$\begin{aligned}
 T_{fi} &= -i \int d^4x A^\mu (-ie) \left[\phi_f^* \partial_\mu \phi_i - \partial_\mu \phi_f^* \phi_i \right] \\
 &= -i \int d^4x A^\mu(x) \underbrace{(-)e N_i N_f (p_i + p_f)_\mu}_{J_\mu^{(fi)}} e^{-i(p_i - p_f)x}
 \end{aligned}$$



2- Matrix element

- Consider the process : $a+1 \rightarrow b+2$



The intermediate field is no more a free particle field but is induced by the motion of the bottom part of the diagram.

The A^μ field is given by the photon propagation equations in presence of a source :

$$(\partial^\nu \partial_\nu) A^\mu(x) - \underbrace{\partial^\mu (\partial^\nu A_\nu)}_{=0 \text{ in Lorentz condition}} = J_{(12)}^\mu$$

- The current is given by :

$$J_\mu^{(12)} = \langle p_2 | \hat{J}_\mu | p_1 \rangle = -eN_1N_2 (p_1 + p_2)_\mu e^{-i(p_1-p_2).x}$$

2. Matrix element

- The previous expression allows to compute the A^μ field :

$$\square A_\mu = J_\mu^{(12)} = -eN_1N_2 (p_1 + p_2)_\mu e^{-i(p_1-p_2)\cdot x} = -eN_1N_2 (p_1 + p_2)_\mu e^{iq\cdot x}$$

$$\Rightarrow A_\mu = \langle 0 | \hat{A}_\mu | q \rangle = +\frac{1}{q^2} eN_1N_2 (p_1 + p_2)_\mu e^{iq\cdot x}$$

$$\Rightarrow A_\mu = \frac{-g_\mu^\nu}{q^2} J_\nu^{(12)}$$

- Replacing this expression in the matrix element :

$$T_{fi} = -i \int d^4x J_\mu^{(ab)} \frac{-g^{\mu\nu}}{q^2} J_\nu^{(12)}$$

- Namely :


$$T_{fi} = -i(-e)(-e)N_aN_bN_1N_2 (p_a + p_b)_\mu \frac{-g^{\mu\nu}}{q^2} (p_1 + p_2)_\nu$$

$$\times (2\pi)^4 \delta^4(p_f - p_i - k)$$

3- Feynman rules for QED

Summary : “reading” a Feynman diagram

- The basic elements of a diagram are :
 - the external lines
 - the vertex operator
 - the internal lines or propagators



with the transferred momentum : $q = (p_a - p_b) = (p_2 - p_1)$

3- Feynman rules for QED

Including fermions into the game: same procedure

⇒ matrix element reduced expression

$$T_{fi} = -i \int d^4x J_{\mu}^{(ab)} \frac{-g^{\mu\nu}}{q^2} J_{\nu}^{(12)}$$

⇒ with a current given by (see lecture on Dirac's equation) :

$$J_{\mu}^{(fi)} = -e (\bar{\psi}_f \gamma_{\mu} \psi_i) = -e (\bar{u}_f \gamma_{\mu} u_i) e^{i(p_f - p_i) \cdot x}$$

⇒ N.B. to be compared with the spinless case :

$$J_{\mu}^{(fi)} = -e (p_f + p_i)_{\mu} e^{i(p_f - p_i) \cdot x}$$

3- Feynman rules for QED

Feynman rules for QED:

- External lines:

- fermions => $\xrightarrow{\text{(incoming)}} \bigcirc$
 $u(p, s)$

$\bigcirc \xrightarrow{\text{(outgoing)}}$
 $\bar{u}(p', s')$

- anti-fermions => $\xrightarrow{\text{(incoming)}} \bigcirc$
 $\bar{v}(p, s)$

$\bigcirc \xrightarrow{\text{(outgoing)}}$
 $v(p', s')$

- spin-0 bosons => $\xrightarrow{\text{(incoming)}} \bigcirc$
 cst

$\bigcirc \xrightarrow{\text{(outgoing)}}$
 cst

- photons => $\xrightarrow{\text{(incoming)}} \bigcirc$
 $\epsilon_{\mu}(k, \lambda)$

$\bigcirc \xrightarrow{\text{(outgoing)}}$
 $\epsilon_{\mu}^*(k', \lambda')$

3- Feynman rules for QED

Feynman rules for QED:


- Propagators:

- fermions =>



$$\frac{i}{p - m} = \frac{i(p + m)}{p^2 - m^2}$$

- spin-0 bosons =>



$$\frac{i}{p^2 - m^2}$$

- photons =>




$$\frac{-ig^{\mu\nu}}{p^2}$$

3- Feynman rules for QED

Feynman rules for QED:

- Vertices :

- fermions – photon =>



- spin-0 bosons – photon =>

