

---

Travaux Dirigés de Physique Nucléaire

---

Liquid drop model : binding energy, isobaric chains

---

**Exercise 1 :** Energy balance

1 / Find the inequalities between the masses of the neutral "father" and "son" atoms in the case of :

- a  $\beta^-$  decay ?
- a  $\beta^+$  decay ?
- an electron capture by the K level ?

2 / What would be the experimental signatures ?

**Exercise 2 :** Binding energy per nucleon

Starting from the empirical spherical mass formula and neglecting the pairing term, show that the binding energy per particle ( $B/A$ ) for symmetric nuclei  $N = Z = A/2$  has a maximum close to the iron ( $Z = 26$ ). Estimate the value at the maximum.

Numerical values :  $u_s = 18 \text{ MeV}$ ,  $u_c = 0.7 \text{ MeV}$  and  $u_v = 15.8 \text{ MeV}$ .

**Exercise 3 :** Nucleus stability

1 / One considers a nucleus  ${}^A_Z X$  with  $A \gg 1$  and  $Z \gg 1$ . Express the  $Q$ -values of the following decays with respect to the binding energy and its derivatives :

- neutron emission
- proton emission
- $\alpha$ -particle emission
- symmetric fission ( $A \rightarrow A/2$  and  $Z \rightarrow Z/2$ )

**2 /** Deduce the stability of a  ${}_{92}^{238}\text{U}$  nucleus w.r.t. those various decay modes and compute the released energy when the decay is possible. The pairing term will be neglected. Numerical values :  $u_s = 16.8\text{MeV}$ ,  $u_v = 15.5\text{MeV}$ ,  $u_T = 19\text{MeV}$ ,  $u_c = 0.72\text{MeV}$  and  $B_\alpha = 28\text{MeV}$ .

**Exercise 4 :** Neutron stars

**1 /** Extrapolate the mass formula by adding a gravitational potential energy term to study the stability of a neutron star. The surface and pairing terms will be neglected w.r.t. the volume term.

**2 /** What conditions should fulfil the radius and the mass of the neutrons star? More accurate calculations give a limit mass of  $0.1 M_\odot$ . Comment. ( $m_n c^2 = 1.67 \cdot 10^{-27}\text{kg}$  and  $G = 6.673 \cdot 10^{-11} \text{m}^3 \text{kg}^{-1} \text{s}^{-2}$ ,  $M_\odot \sim 2 \cdot 10^{30} \text{kg}$ ).

**Exercise 5 :** Odd isobaric chain

For the isobaric chain  $A = 151$  one gets the following mass excesses :

Z	Element	Mass excess
58	Ce	-62.620
59	Pr	-67.160
60	Nd	-70.957
61	Pm	-73.400
62	Sm	-74.587
63	Eu	-74.663
64	Gd	-74.198
65	Tb	-71.632
66	Dy	-68.902
67	Ho	-63.803
68	Er	-58.500
69	Tm	-51.000
70	Yb	-41.100
71	Lu	-29.010

A	Z	Element	Mass excess
1	1	H	+7.289
4	2	He	+2.424
147	69	Tm	-34.820
150	70	Yb	-37.470

- 1 / Identify the stable isobar(s) of that serie.
- 2 / Give the decay modes of the unstable isobar(s). Summarize the results on a  $M(Z)$  scheme.
- 3 / For  $^{151}\text{Lu}$  indicate all possible decay modes.
- 4 / In which  $Z$  domain should one look for spontaneous neutron emission in that isobaric chain ?

**Exercise 6 : Even isobaric chain**

The following nuclei belong to the  $A = 114$  isobaric chain :



where

- ${}_{46}\text{Pd}$  is a  $\beta^-$  emitter with 1.4 MeV maximal energy of the electrons ;
- ${}_{47}\text{Ag}$  gives a  $\beta^-$  of 4.6 MeV maximal energy, followed by a gamma emission of 0.56 MeV ;
- ${}_{51}\text{Sb}$  gives two  $\beta^+$ , the first one of 2.7 MeV maximal energy followed by a two-gammas cascade of 0.9 MeV and 1.3 MeV, the second one with 3.6 MeV maximum energy followed by a 1.3 MeV gamma.

Find for that particular chain :

- 1 / the coefficients of the mass formula (parabolic form),
- 2 / the stable nucleus (nuclei),
- 3 / the various decay modes of unstable isobar(s) and the maximum energy of the emitted  $\beta$ .