

**A PRECISE DETERMINATION OF THE NUMBER OF FAMILIES WITH LIGHT NEUTRINOS
AND OF THE Z BOSON PARTIAL WIDTHS**

ALEPH Collaboration

D. DECAMP, B. DESCHIZEAUX, J.-P. LEES, M.-N. MINARD

*Laboratoire de Physique des Particules (LAPP), IN2P3-CNRS, F-74019 Annecy-le-Vieux Cedex, France*J.M. CRESPO, M. DELFINO, E. FERNANDEZ ¹, M. MARTINEZ, R. MIQUEL, L.I.M. MIR,
S. ORTEU, A. PACHECO, J.A. PERLAS, E. TUBAU*Laboratorio de Fisica de Altas Energias, Universidad Autonoma de Barcelona, E-08193 Bellaterra (Barcelona), Spain* ²M.G. CATANESI, M. DE PALMA, A. FARILLA, G. IASELLI, G. MAGGI, S. NATALI, S. NUZZO,
A. RANIERI, G. RASO, F. ROMANO, F. RUGGIERI, G. SELVAGGI, L. SILVESTRIS,
P. TEMPESTA, G. ZITO*INFN, Sezione di Bari e Dipartimento di Fisica dell' Università, I-70126 Bari, Italy*H. HU, D. HUANG, J. LIN, T. RUAN, T. WANG, W. WU, Y. XIE, D. XU, R. XU, J. ZHANG,
W. ZHAO*Institute of High-Energy Physics, Academia Sinica, Beijing, P.R. China* ³H. ALBRECHT ⁴, W.B. ATWOOD ⁵, F. BIRD, E. BLUCHER, T.H. BURNETT ⁶, T. CHARITY,
H. DREVERMANN, LI. GARRIDO, C. GRAB, R. HAGELBERG, S. HAYWOOD, B. JOST,
M. KASEMANN, G. KELLNER, J. KNOBLOCH, A. LACOURT, I. LEHRAUS, T. LOHSE,
D. LÜKE ⁴, A. MARCHIORO, P. MATO, J. MAY, A. MINTEN, A. MIOTTO, P. PALAZZI,
M. PEPE-ALTARELLI, F. RANJARD, A. ROTH, J. ROTHBERG ⁶, H. ROTSCHEIDT,
W. VON RÜDEN, R. ST.DENIS, D. SCHLATTER, M. TAKASHIMA, M. TALBY, H. TAUREG,
W. TEJESSY, H. WACHSMUTH, S. WHEELER, W. WIEDENMANN, W. WITZELING,
J. WOTSCHACK*European Laboratory for Particle Physics (CERN), CH-1211 Geneva 23, Switzerland*Z. AJALTOUNI, M. BARDADIN-OTWINOWSKA, A. FALVARD, P. GAY, P. HENRARD,
J. JOUSSET, B. MICHEL, J-C. MONTRET, D. PALLIN, P. PERRET, J. PRORIOL, F. PRULHIÈRE
*Laboratoire de Physique Corpusculaire, Université Blaise Pascal, IN2P3-CNRS, Clermont-Ferrand, F-63177 Aubière, France*J.D. HANSEN, J.R. HANSEN, P.H. HANSEN, R. MØLLERUD, G. PETERSEN
Niels Bohr Institute, DK-2100 Copenhagen, Denmark ⁷

E. SIMOPOULOU, A. VAYAKI

Nuclear Research Center Demokritos (NRCD), Athens, Greece

J. BADIER, A. BLONDEL, G. BONNEAUD, J. BOUROTE, F. BRAEMS, J.C. BRIENT,
 M.A. CIOCCI, G. FOUQUE, R. GUIREL, A. ROUGÉ, M. RUMPF, R. TANAKA, H. VIDEAU,
 I. VIDEAU¹

Laboratoire de Physique Nucléaire et des Hautes Energies, Ecole Polytechnique, IN2P3-CNRS, F-91128 Palaiseau Cedex, France

D.J. CANDLIN

*Department of Physics, University of Edinburgh, Edinburgh EH9 3JZ, UK*⁸

A. CONTI, G. PARRINI

Dipartimento di Fisica, Università di Firenze, INFN Sezione di Firenze, I-50125 Florence, Italy

M. CORDEN, C. GEORGIOPoulos, J.H. GOLDMAN, M. IKEDA, J. LANNUTTI,
 D. LEVINTHAL⁹, M. MERMIKIDES, L. SAWYER, G. STIMPFL

*Supercomputer Computations Research Institute and Department of Physics, Florida State University,
 Tallahassee, FL 32306, USA*^{10,11,12}

A. ANTONELLI, R. BALDINI, G. BENCIVENNI, G. BOLOGNA¹³, F. BOSSI, P. CAMPANA,
 G. CAPON, V. CHIARELLA, G. DE NINNO, B. D'ETTORRE-PIAZZOLI¹⁴, G. FELICI,
 P. LAURELLI, G. MANNOCCHI¹⁴, F. MURTAS, G.P. MURTAS, G. NICOLETTI, P. PICCHI¹³,
 P. ZOGRAFOU

Laboratori Nazionali dell' INFN (LNF-INFN), I-00044 Frascati, Italy

B. ALTOON, O. BOYLE, A.W. HALLEY, I. TEN HAVE, J.L. HEARNS, I.S. HUGHES, J.G. LYNCH,
 W.T. MORTON, C. RAINES, J.M. SCARR, K. SMITH¹, A.S. THOMPSON

*Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK*⁸

B. BRANDL, O. BRAUN, R. GEIGES, C. GEWENIGER¹, P. HANKE, V. HEPP, E.E. KLUGE,
 Y. MAUMARY, M. PANTER, A. PUTZER, B. RENSCH, A. STAHL, K. TITTEL, M. WUNSCH
*Institut für Hochenergiephysik, Universität Heidelberg, D-6900 Heidelberg, FRG*¹⁵

A.T. BELK, R. BEUSELINCK, D.M. BINNIE, W. CAMERON¹, M. CATTANEO, P.J. DORNAN,
 S. DUGEAY, R.W. FORTY, A.M. GREENE, J.F. HASSARD, S.J. PATTON, J.K. SEDGBEER,
 G. TAYLOR, I.R. TOMALIN, A.G. WRIGHT

*Department of Physics, Imperial College, London SW7 2BZ, UK*⁸

P. GIRTNER, D. KUHN, G. RUDOLPH

*Institut für Experimentalphysik, Universität Innsbruck, A-6020 Innsbruck, Austria*¹⁶

C.K. BOWDERY¹, T.J. BRODBECK, A.J. FINCH, F. FOSTER, G. HUGHES, N.R. KEEMER,
 M. NUTTALL, B.S. ROWLINGSON, T. SLOAN, S.W. SNOW

*Department of Physics, University of Lancaster, Lancaster LA1 4YB, UK*⁸

T. BARCZEWSKI, L.A.T. BAUERDICK, K. KLEINKNECHT¹, B. RENK, S. ROEHN,
 H.-G. SANDER, M. SCHMELLING, F. STEEG

*Institut für Physik, Universität Mainz, D-6500 Mainz, FRG*¹⁵

J.-P. ALBANESE, J.-J. AUBERT, C. BENCHOUK, A. BONISSENT, D. COURVOISIER,
 F. ETIENNE, E. MATSINOS, S. PAPALEXIOU, P. PAYRE, B. PIETRZYK¹, Z. QIAN
Centre de Physique des Particules, Faculté des Sciences de Luminy, IN2P3-CNRS, F-13288 Marseille, France

W. BLUM, P. CATTANEO, G. COWAN, B. DEHNING, H. DIETL, M. FERNANDEZ-BOSMAN,
 D. HAUFF, A. JAHN, E. LANGE, G. LÜTJENS, G. LUTZ, W. MÄNNER, H.-G. MOSER, Y. PAN,
 R. RICHTER, A.S. SCHWARZ, R. SETTLES, U. STIEGLER, U. STIERLIN, J. THOMAS,
 G. WALTERMANN

*Max-Planck-Institut für Physik und Astrophysik, Werner-Heisenberg-Institut für Physik, D-8000 Munich, FRG*¹⁵

V. BERTIN, G. DE BOUARD, J. BOUCROT, O. CALLOT, X. CHEN, A. CORDIER, M. DAVIER,
 G. GANIS, J.-F. GRIVAZ, Ph. HEUSSE, P. JANOT, V. JOURNÉ, D.W. KIM, J. LEFRANÇOIS,
 A.-M. LUTZ, J.-J. VEILLET, F. ZOMER

Laboratoire de l'Accélérateur Linéaire, Université de Paris-Sud, IN2P3-CNRS, F-91405 Orsay Cedex, France

S.R. AMENDOLIA, G. BAGLIESI, G. BATIGNANI, L. BOSISIO, U. BOTTIGLI, C. BRADASCHIA,
 I. FERRANTE, F. FIDECARO, L. FOÀ¹, E. FOCARDI, F. FORTI, A. GIASSI, M.A. GIORGI,
 F. LIGABUE, A. LUSIANI, E.B. MANNELLI, P.S. MARROCCHESI, A. MESSINEO, F. PALLA,
 G. SANGUINETTI, S. SCAPELLATO, J. STEINBERGER, R. TENCHINI, G. TONELLI,
 G. TRIGGIANI

Dipartimento di Fisica dell' Università, INFN Sezione di Pisa, e Scuola Normale Superiore, I-56010 Pisa, Italy

J.M. CARTER, M.G. GREEN, P.V. MARCH, T. MEDCALF, M.R. SAICH, J.A. STRONG¹,
 R.M. THOMAS, T. WILDISH

*Department of Physics, Royal Holloway & Bedford New College, University of London, Surrey TW20 0EX, UK*⁸

D.R. BOTTERILL, R.W. CLIFFT, T.R. EDGECOCK, M. EDWARDS, S.M. FISHER, J. HARVEY,
 T.J. JONES, P.R. NORTON, D.P. SALMON, J.C. THOMPSON

*Particle Physics Department, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, UK*⁸

E. AUBOURG, B. BLOCH-DEVAUX, P. COLAS, C. KLOPFENSTEIN, E. LANÇON, E. LOCCI,
 S. LOUCATOS, L. MIRABITO, E. MONNIER, P. PEREZ, F. PERRIER, J. RANDER,
 J.-F. RENARDY, A. ROUSSARIE, J.-P. SCHULLER

*Département de Physique des Particules Élémentaires, CEN-Saclay, F-91191 Gif-sur-Yvette Cedex, France*¹⁷

J.G. ASHMAN, C.N. BOOTH, F. COMBLEY, M. DINSDALE, J. MARTIN, D. PARKER,
 L.F. THOMPSON

*Department of Physics, University of Sheffield, Sheffield S3 7RH, UK*⁸

S. BRANDT, H. BURKHARDT, C. GRUPEN, H. MEINHARD, E. NEUGEBAUER, U. SCHÄFER,
 H. SEYWERD

*Fachbereich Physik, Universität Siegen, D-5900 Siegen, FRG*¹⁵

B. GOBBO, F. LIELLO, E. MILOTTI, F. RAGUSA¹⁸, L. ROLANDI¹

Dipartimento di Fisica, Università di Trieste e INFN Sezione di Trieste, I-34127 Trieste, Italy

L. BELLANTONI, J.F. BOUDREAU, D. CINABRO, J.S. CONWAY, D.F. COWEN, Z. FENG, J.L. HARTON, J. HILGART, R.C. JARED ¹⁹, R.P. JOHNSON, B.W. LECLAIRE, Y.B. PAN, T. PARKER, J.R. PATER, Y. SAADI, V. SHARMA, J.A. WEAR, F.V. WEBER, SAU LAN WU, S.T. XUE and G. ZOBERNIG

Department of Physics, University of Wisconsin, Madison, WI 53706, USA ²⁰

Received 19 December 1989

More extensive and precise results are reported on the parameters of Z decay. On the basis of 20 000 Z decays collected with the ALEPH detector at LEP we find $M_Z = 91.182 \pm 0.026$ (exp.) ± 0.030 (beam) GeV, $\Gamma_Z = 2.541 \pm 0.056$ GeV and $\sigma_{\text{had}}^0 = 41.4 \pm 0.8$ nb. The partial widths for the hadronic and leptonic channels are $\Gamma_{\text{had}} = 1804 \pm 44$ MeV, $\Gamma_{e^+e^-} = 82.1 \pm 3.4$ MeV, $\Gamma_{\mu^+\mu^-} = 87.9 \pm 6.0$ MeV and $\Gamma_{\tau^+\tau^-} = 86.1 \pm 5.6$ MeV, in good agreement with the standard model. On the basis of the average leptonic width $\Gamma_{e^+e^-} = 83.9 \pm 2.2$ MeV, the effective weak mixing angle is found to be $\sin^2 \theta_w(M_Z) = 0.231 \pm 0.008$. Using the partial widths calculated in the standard model, the number of light neutrino families is $N_\nu = 3.01 \pm 0.15$ (exp.) ± 0.05 (theor.).

1. Introduction

With the start-up of LEP, one of the interesting questions which could be addressed was that of the number of neutrino species into which the Z decays, not by the direct observation of the neutrinos, but by

a precise measurement of the Z width, or of the cross-section at the Z peak. The latter is statistically more powerful, but systematically more demanding because it requires precise knowledge of the absolute luminosity and of the Z decay detection efficiency. Several results based on the cross-section have been reported recently [1–5] and are in agreement with three neutrino families; however, the most precise of these [5] is only $2\frac{1}{2}$ standard deviations away from four families. We therefore believe it useful to report the continuation of our work, based on a six-fold statistical increase in the data sample. The analysis has remained basically the same, but the detector has continuously improved and our understanding of the systematic errors has progressed. In addition to the number of neutrino families N_ν , new results for the Z mass and line shape, as well as the leptonic branching ratios are given. For a brief description of the detector, we refer to ref. [5]. A more detailed description is in preparation [6].

2. Event triggers

Two independent triggers are used in this analysis (i) a basic trigger based on the electromagnetic calorimeter (ECAL), which requires a total energy of 6.5 GeV in the ECAL barrel, 3.8 GeV in either ECAL end-cap, or 1.6 GeV in both end-caps; (ii) a penetrating-charged-particle trigger, which requires that at least 5 inner tracking chamber (ITC) wire planes and

¹ Present address: CERN, CH-1211 Geneva 23, Switzerland.

² Supported by CAICYT, Spain.

³ Supported by the National Science Foundation of China.

⁴ Permanent address: DESY, D-2000 Hamburg 52, FRG.

⁵ On leave of absence from SLAC, Stanford, CA 94309, USA.

⁶ On leave of absence from University of Washington, Seattle, WA 98195, USA.

⁷ Supported by the Danish Natural Science Research Council.

⁸ Supported by the UK Science and Engineering Research Council.

⁹ Supported by SLOAN fellowship, contract BR 2703.

¹⁰ Supported by the US Department of Energy, contract DE-FG05-87ER40319.

¹¹ Supported by the NSF, contract PHY-8451274.

¹² Supported by the US Department of Energy, contract DE-FC05-85ER250000.

¹³ Also at Istituto di Fisica Generale, Università di Torino, Turin, Italy.

¹⁴ Also at Istituto di Cosmo-Geofisica del CNR, Turin, Italy.

¹⁵ Supported by the Bundesministerium für Forschung und Technologie, FRG.

¹⁶ Supported by Fonds zur Förderung der wissenschaftlichen Forschung, Austria.

¹⁷ Supported by the Institut de Recherche Fondamentale du CEA.

¹⁸ Present address: INFN, Sezione di Milano, I-20133 Milan, Italy.

¹⁹ Permanent address: LBL, Berkeley, CA 94720, USA.

²⁰ Supported by the US Department of Energy, contract DE-AC02-76ER00881.

at least 4 hadron calorimeter (HCAL) double planes give signals in the same azimuthal region.

3. Z event selection and efficiency

Events are selected in two independent ways, one based on calorimetric energy, the other on charged tracks. Both require the basic trigger; the other trigger is used exclusively to measure the trigger efficiency.

The calorimetric selection requires a minimum total electromagnetic and hadronic energy of at least 20 GeV. In addition it is required that at least 6 GeV of this energy be in the ECAL barrel or 1.5 GeV in each of the ECAL end-caps. If there are at least 5 time projection chamber (TPC) tracks (95% of the events) there are no further requirements. For events with 1 to 4 tracks, additional cuts are imposed to eliminate e^+e^- and $\mu^+\mu^-$ final states. Bhabha's are rejected on the basis of their characteristic tight energy clusters in the electromagnetic calorimeter. To eliminate muonic decays, events are rejected if there are exactly two tracks, each with associated hits (4 or more strips in the 10 outer layers) in HCAL. For events with no tracks at all, two additional cuts are applied to eliminate cosmic rays: a time window of ± 350 ns based on timing in ECAL, and a requirement of at least two clusters of 3 GeV each in ECAL. This selection leaves 18 654 events: 16 930 with 5 or more tracks, 872 with 1 to 4 tracks, and 852 with no tracks. This latter class includes 804 events with the TPC turned off. All events with less than 5 tracks were visually inspected: 121 events were identified as Bhabha events in which one or both electrons are in the cracks of ECAL but the energy could be seen in HCAL, 14 events are classified as beam-gas interactions, 3 as muon pairs, and 92 could not be identified, so that 46 with an error of ± 46 were included in the final event sample. As a check, in the visual inspection, $\tau^+\tau^-$ decays were identified. Almost all $\tau^+\tau^-$ events should appear in the 1-4 track category. On the basis of the branching ratio $\Gamma_{\tau\tau}/\Gamma_{\text{had}} = 0.048$ and the Monte Carlo simulation, 510 $\tau\tau$ events are expected, and 524 were found. This identification, however, is not used in the analysis.

We are left with a total of 18 470 events. The distribution in total calorimetric energy is shown in fig. 1a, together with the Monte Carlo prediction. Fig. 1b




Fig. 1. Distributions for calorimetrically selected events. (a) Total calorimetric energy; (b) absolute value of the cosine of the thrust axis.

shows the distribution of the calorimetrically determined thrust axis. The experimental energy distribution is somewhat wider than the simulation. This difference is due to noise problems in the hadron calorimeter but has a negligible effect on the detection efficiency; the only use made here of HCAL is the 20 GeV total calorimetric energy requirement. This

conclusion is corroborated by the agreement between data and Monte Carlo at low energy in fig. 1a. The calculated efficiencies are $(99.4 \pm 0.2)\%$ for hadronic events, and $(60 \pm 6)\%$ for $\tau^+\tau^-$ events. The possible contamination by the two photon process ($\gamma\gamma$) is estimated using the sample at the lowest energy, in which this background would be concentrated. We study the cross-section of the data with calorimetric energy between 20 and 35 GeV (330 events) as a function of the beam energy. The Z component must exhibit the resonance characteristics whereas the $\gamma\gamma$ component is almost independent of energy. There is no indication of a $\gamma\gamma$ component. The upper limit for the whole data sample due to $\gamma\gamma$ background is 0.1%. The total systematic uncertainty in the efficiency for this selection is estimated to be 0.6%.

The charged track selection requires at least 5 tracks in the TPC with the sum of the track energies (assuming the pion mass) larger than 10% of the center-of-mass energy. The tracks must have a polar angle above 18.2° . This ensures that at least 6 TPC pad rows are traversed; at least 4 reconstructed coordinates per track are required. The distance of closest approach of the reconstructed tracks from the collision point must be less than 10 cm along the beam and 2 cm transverse to it. Track reconstruction efficiency is $\sim 99\%$. Distributions based on the tracks, such as multiplicity, total energy, sphericity, thrust, etc. are in good agreement [5,7] with hadronization models [8]. For the accepted events, distributions of the total energy and sphericity axis are shown in fig. 2a and 2b, respectively. The calculated efficiency for $q\bar{q}$ events is $(97.5 \pm 0.6)\%$. The background of $\tau^+\tau^-$ events is estimated at (30 ± 10) events, and is subtracted. The background from the two-photon process is calculated to be 15 pb. To check the two-photon contribution, the cross-section for events with track energy between $0.1E_{CM}$ and $0.15E_{CM}$ is studied as a function of center of mass energy to separate the resonant and non-resonant contributions. A non-resonant cross-section of (-2 ± 32) pb is found, and (15 ± 15) pb is subtracted from the observed hadronic cross-section. The total systematic uncertainty in the efficiency for the track selected sample is also estimated at 0.6%.

The trigger efficiency is measured by means of the penetrating track trigger, which is entirely independent, and which overlapped in 92% of the events.




Fig. 2. Distributions for track selected events. (a) Total track energy; (b) absolute value of the cosine of the sphericity axis.

Events triggered by the penetrating track trigger and missed by the basic trigger correspond to 0.05% of the accepted event sample. Conservatively, the trigger inefficiency is $(0.1 \pm 0.1)\%$ for both event samples.

4. Luminosity detector, acceptance conditions and trigger

Accurate measurement of the luminosity is essential to the precise determination of the peak cross-section especially because the acceptance for the Z events is very close to unity and the uncertainty correspondingly small. The Bhabha events used to measure the luminosity have a polar angle dependence of $\sin^{-4}(\frac{1}{2}\theta)$; a clear and well understood acceptance at the inner boundary is therefore crucial.

The ALEPH luminosity detector consists of a wire chamber arrangement followed by a calorimeter (LCAL) around the beam pipe at small angles on both sides of the detector. Only the calorimeter is used in the present analysis. It is a 38 layer lead and proportional-wire chamber sandwich construction, read out in projective towers, 784 on each side (see fig. 3a). Each tower is read out in 3 stories, 4.7, 10.4 and 9.4 radiation lengths in depth, respectively. There are two semicylindrical modules on each side covering the polar angle range 42–160 mrad. Within each module, the mechanical precision in the relative position of the towers is 120 μm . The uncertainty on the inner radius due to the relative positioning of the two modules is 140 μm . The distance of the shower maximum from the collision point is 280 cm. The energy resolution is 3%, tower-to-tower response uniformity $\pm 2\%$, and the spatial resolution of the shower position 1.2 mm in x and y .

The acceptance region is defined by a boundary between towers as shown for one half of one module in fig. 3a. The acceptance region excludes an inner band of towers and further excludes a large outer region, essentially the region outside ~ 110 mrad. The latter exclusion results in a loss of $\sim 15\%$ of the luminosity events, but was desirable, since this region is shadowed by the end-plate of the ITC and TPC, degrading the electron energies somewhat and consequently introducing a systematic error. Acceptance requires that the electron or positron shower pulse height in the front storey is larger in the acceptance region than in the boundary region on one side. There is no such requirement on the opposite side other than that the shower position be inside a 125 mrad contour. The side on which the boundary conditions must be met alternates from one event to the next. The asymmetry in the acceptance condition, together with the re-

quirement of alternation from event to event, ensure that the acceptance is independent, in first order, of transverse and longitudinal displacement of the collision point and of small displacements of the beam direction. The pulse height requirement across the fiducial boundary results in a remarkably precise definition of the fiducial region: the position uncertainty at the boundary is $\sim 150 \mu\text{m}$. The projective nature of the geometry is essential here. The x - y position of the beam crossing relative to the calorimeters is measured for each LEP fill on the basis of the shower positions on the two sides with a precision of ~ 0.2 mm, but the consequent corrections are negligible.

In addition to the geometric requirement, it is required that the energy on each side is greater than $0.22E_{\text{CM}}$, and the sum greater than $0.6E_{\text{CM}}$.

Three triggers were used in the luminosity determination:

(i) A basic luminosity trigger. This trigger requires a coincidence between the two sides with more than 20 GeV on one and 16 GeV on the other.

(ii) A high level single trigger, requiring more than 31 GeV on either side. This trigger serves as a check and permits the efficiency determination of the basic trigger.

(iii) Prescaled single triggers of more than 20 GeV and 16 GeV on either side. These triggers enable determination of the background due to random coincidences of off-momentum electrons.

The initial data sample [5] had a small problem in the coincidence trigger, which has since been corrected. The luminosity trigger inefficiency for the total sample is measured to be $(0.1 \pm 0.1)\%$.

Fig. 3b shows the polar-angle distribution of accepted events on the fiducial side, for data and simulation. Fig. 3c shows the total energy distribution. Fig. 3d shows the distribution in the difference, $\Delta\phi$, between the azimuthal angles in the two arms after correction for beam position and magnetic deflection. The Bhabha events are near 180° ; a cut at 170° is applied. The data shown in this figure also permit an estimate of the background due to random coincidences in the two sides between single, off-momentum particles. This background is distributed much more broadly in $\Delta\phi$ than the luminosity events. The events in the region $0^\circ < \Delta\phi < 10^\circ$ and $160^\circ < \Delta\phi < 170^\circ$ can be used to subtract the background at




Fig. 3. The luminosity calorimeter. (a) Tower structure in one quadrant showing acceptance boundary; (b) distribution of the polar angle; (c) distribution of the total Bhabha energy relative to the collision energy, before and after background subtraction. The minimum value for acceptance is 0.6; (d) distribution of the difference in the azimuthal angle between the two sides. The acceptance region is $170^\circ < \Delta\phi < 180^\circ$.

$170^\circ < \Delta\phi < 180^\circ$. As can be seen in fig. 3d, the background is small, at a level of $\sim 0.4\%$.

The effective cross-section is calculated using an event generator that includes first-order radiative corrections [9]. The events were generated at 91.0

GeV collision energy and the cross-section is found to be 26.78 ± 0.15 (stat.) ± 0.20 (theor.) nb at 91.0 GeV. For other energies this cross-section was multiplied by the factor $(91 \text{ GeV}/E_{\text{CM}})^2$ and corrected for small ($< 1\%$) electroweak effects.

We checked that the cross-section within our cuts is not very sensitive to radiative corrections. The difference between the lowest-order and the first-order calculations is about 1%. The systematic error introduced by neglecting higher orders is expected to be smaller than this value; we assume an error of 0.7%. The hadronic vacuum polarization has been included [10]. The uncertainty for the small momentum transfer involved is of order 10^{-4} , and is therefore negligible.

The systematic relative errors in the determination of the luminosity are estimated as shown in table 1.

Some checks on the systematic quality of the luminosity measurements were possible. The precision of the geometrical event selection could be checked by repeating the selection, but basing it not on the first calorimetric layer of 4.7 radiation lengths, but instead on the second of 10.4 radiation lengths which contains the bulk of the shower. In a sub-sample of the data, 4169 events were selected on the basis of the first layer and 4171 on the basis of the second. A total of 6 events were not in common. A second, more comprehensive, check was possible by comparing the result to that obtained on the basis of a more restrictive selection in which the inner boundary is displaced by one tower width. The ratio of accepted events was found to be 0.732 ± 0.004 compared to the Monte Carlo expectation of 0.729. In a third check, the energy requirement was changed from the fraction of 0.6 for the total energy to 0.66 and 0.55 of the beam energy for the showers on the fiducial and non-fiducial sides respectively. The acceptance changed by 1.5% compared to the Monte Carlo expectation of 1%. The discrepancy of 0.5% is due to known flaws in the detector simulation, but is below the given error of 0.9%.

5. Cross-sections on the Z resonance

Data were obtained at 11 energies: at the Z peak, and at the peak ± 0.25 GeV, ± 1.0 GeV, ± 1.5 GeV, ± 2.0 GeV, ± 3.0 GeV and $+4.0$ GeV. The results reported here are based on $\sim 20\,000$ Z decays which include the 3300 already reported [5]. The calorimetric selection did not require the TPC, so that its integrated luminosity is higher than that for the track selection. For the data accessible to both selections, 96% of the events are in common. The remaining 4% are found only calorimetrically: $\frac{2}{3}$ of these are $\tau^+\tau^-$ decays and the remaining $\frac{1}{3}$ are hadronic decays. These differences are as expected from the simulation. A detailed comparison of the two selections for the events with at least 5 tracks finds an agreement of better than 0.3%.

The number of Z and luminosity events are given in table 2 for the different energies and the two event selections. The resultant hadronic cross-sections are listed as well. In the case of the track selection,

$$\sigma_{\text{had}} = \frac{N_{\text{track}}}{\epsilon_{\text{track}}} \frac{\sigma_{\text{Lum}}}{N_{\text{Lum}}},$$

where

$$\epsilon_{\text{track}} = 0.975 \pm 0.006.$$

In the case of calorimetric selection,

$$\sigma_{\text{had}} = \frac{N_{\text{cal}}}{\epsilon_{\text{cal}}} \frac{\sigma_{\text{Lum}}}{N_{\text{Lum}}},$$

where

$$\begin{aligned} \epsilon_{\text{cal}} &= \epsilon_{\text{cal, had}} + \frac{\sigma_{\tau}}{\sigma_{\text{had}}} \epsilon_{\text{cal, } \tau} \\ &= 0.994 + 0.048 \times 0.60 \\ &= 1.022 \pm 0.006 \text{ (at the peak),} \end{aligned}$$

Table 1

position of pad towers within a module and between the two halves	0.002
inadequacy of simulation	0.009
energy resolution and cell to cell variation	0.003
uncertainty in collision position and beam angle relative to calorimeters	negligible
theoretical uncertainty, higher order radiative effects	0.007
statistics of simulation	0.004
total estimated systematic error	0.013

Table 2

Results for σ_{had} near the Z peak. The number of Z decay candidates and luminosity event numbers and resultant hadronic cross-sections for the two event selections, as well as the combined result are shown. The errors are statistical only.

Energy (GeV)	Track selection			Calorimeter selection			Average cross-section (nb)
	N_{track}	N_{Lum}	cross-section (nb)	N_{cal}	N_{Lum}	cross-section (nb)	
88.278	306	2015	4.43 ± 0.27	405	2447	4.56 ± 0.25	4.49 ± 0.25
89.289	337	1020	9.45 ± 0.59	546	1570	9.46 ± 0.47	9.46 ± 0.47
90.285	1620	2310	19.65 ± 0.63	1714	2341	19.58 ± 0.62	19.61 ± 0.62
91.036	4057	3691	30.16 ± 0.68	4812	4188	30.15 ± 0.64	30.15 ± 0.64
91.288	4312	3835	30.58 ± 0.68	4579	3874	30.82 ± 0.68	30.70 ± 0.68
91.531	2884	2680	29.04 ± 0.78	3501	3136	28.79 ± 0.71	28.92 ± 0.71
92.286	1340	1686	21.01 ± 0.78	1429	1715	21.00 ± 0.76	21.00 ± 0.76
92.565	130	222	15.40 ± 1.71	143	225	15.91 ± 1.73	15.65 ± 1.71
93.292	827	1585	13.50 ± 0.58	913	1685	13.36 ± 0.56	13.43 ± 0.56
94.278	234	688	8.62 ± 0.66	258	715	8.61 ± 0.64	8.61 ± 0.64
95.036	95	396	5.98 ± 0.69	170	662	6.00 ± 0.53	5.99 ± 0.53

where the error includes all systematic uncertainties.

In the following analysis, the average result of the two selections, also given in table 2, is used. The systematic errors are not included in table 2. They are 1.3% for the luminosity measurement, 0.6% for the Z events of each selection, and 0.4% for the corresponding error in the combined cross-section.

6. The resonance parameters

The energy dependence of the cross-section near the Z resonance is expected to have the Breit-Wigner form:

$$\sigma_{\text{had}} = \sigma_{\text{had}}^0 \frac{s\Gamma_Z^2}{(s - M_Z^2)^2 + s^2\Gamma_Z^2/M_Z^2} [1 + \delta_{\text{rad}}(s)],$$

$$\sigma_{\text{had}}^0 = 12\pi \frac{\Gamma_{ee}\Gamma_{\text{had}}}{M_Z^2\Gamma_Z^2}. \quad (1)$$

Here, Γ_{ee} and Γ_{had} are the partial widths for Z decay to e^+e^- and $q\bar{q}$, respectively, Γ_Z is the total Z width, and M_Z is the Z mass. To obtain a meaningful cross-section the bremsstrahlung from the initial state has to be resummed to all orders as described in refs. [11–15]. The effect of these corrections is summarized in the term $\delta_{\text{rad}}(E)$ of (1), which, although of the order of 30% at the peak, is known to better than 0.5%.

A three-parameter fit to the data on the basis of eq. (1) and the computer program of Burgers [15] and

the formula of Borelli et al. [16] yields the peak cross-section σ_{had}^0 , the Z mass and Γ_Z :

$$\sigma_{\text{had}}^0 = 41.4 \pm 0.8 \text{ nb},$$

$$M_Z = 91.182 \pm 0.026 \text{ (exp.)} \pm 0.030 \text{ (beam) GeV},$$

$$\Gamma_Z = 2.541 \pm 0.056 \text{ GeV}.$$

The beam error in M_Z is due to the uncertainty in the mean e^+e^- collision energy. This error was determined by the LEP Division [17].

The absolute scale error from luminosity and acceptance systematic errors introduces correlations between the points, and these correlations have been taken into account in the fit. In the result for M_Z , the error due to the uncertainty in the collision energy is stated separately. The possible changes in the beam energy from one run period to another do not contribute significantly to any of these quantities.

The χ^2 of the fit is 10.5 for 8 degrees of freedom. The cross-sections and the expectations for two, three or four neutrinos are presented in fig. 4. Fig. 5 shows the 68% and 99% confidence limit contours in the $\sigma_{\text{had}}^0 - \Gamma_Z$ plane. The results are in agreement with, but considerably more precise than, the measurements of only two months ago [1–5].

So far this determination of the Z resonance parameters has been essentially model independent. It is possible to use the standard model predictions for the partial widths [18],




Fig. 4. Cross-section for $e^+e^- \rightarrow$ hadrons as function of LEP energy. The standard model predictions for $N_\nu=2, 3$ and 4 are shown.




Fig. 5. Contours of constant χ^2 in the $\sigma^0_{\text{had}}-\Gamma_Z$ plane, and the standard model predictions for $N_\nu=2, 3$ and 4.

$$\Gamma_{\text{had}} = 1737 \pm 22 \text{ MeV},$$

$$\Gamma_{\ell^+\ell^-} = 83.5 \pm 0.5 \text{ MeV}$$

$$\Gamma_{\nu\nu} = 166.5 \pm 1.0 \text{ MeV},$$

together with $\Gamma_Z = \Gamma_{\text{had}} + 3\Gamma_{\ell^+\ell^-} + N_\nu\Gamma_{\nu\nu}$ to find N_ν on the basis of the measured total width Γ_Z . The result is $N_\nu = 3.30 \pm 0.37$.

7. Number of neutrino families

In the frame of the standard model, the statistically most powerful way to obtain the number of neutrino families is not from the measurement of the width, but from the peak cross-section, σ^0_{had} . The data are refitted with only N_ν and M_Z as parameters. Since the mass determination is independent of the absolute cross-section and the width, M_Z changes only slightly in this fit. We find

$$M_Z = 91.175 \pm 0.027 \pm 0.030 \text{ GeV},$$

$$N_\nu = 3.01 \pm 0.15 \text{ (exp.)} \pm 0.05 \text{ (theor.)}.$$

The theoretical uncertainty is due to uncertainties in the partial width calculation and is discussed in our previous letter [5]. The χ^2 of this fit is 11.3 for 9 degrees of freedom. The agreement with the earlier results [1–5], which give an average of $N_\nu = 3.13 \pm 0.25$, is good.

8. Leptonic branching ratios

The leptonic branching ratio of the Z [19,20] has been determined, using the larger data sample, in a way similar to that reported in ref. [19]. The result of the detailed analysis is given in table 3. The e^+e^- and $\tau^+\tau^-$ channels were analyzed in a manner identical to ref. [19] using the same range of energies. A somewhat larger data sample, however, was used for the $\tau^+\tau^-$ than for the e^+e^- . For the $\mu^+\mu^-$ channel a slightly different procedure to ref. [19] was adopted in order to extend the angular range to $|\cos \theta| < 0.90$ and data at all center-of-mass energies were used. Candidate events for $\mu^+\mu^-$ pairs were selected as in ref. [19]. However, muons were identified here by requiring also that at least one of the two energetic tracks to have a low energy deposit in the ECAL, compatible with a minimum ionizing particle. Some runs in which the HCAL readout was not operational were excluded. The principal backgrounds in each channel in table 3 are lepton pairs mistaken as another type of lepton pair. The last row in table 3 gives

Table 3

Details of the evaluation of the corrected number of events for the three channels $Z \rightarrow \ell^+ \ell^-$. The last row gives the ratio (P) of leptonic events to hadronic events of the same data sample.

	$e^+ e^-$			$\mu^+ \mu^-$			$\tau^+ \tau^-$		
$\ell^+ \ell^-$ candidates	398	\pm	20				382	\pm	20
background events	14						2	\pm	1
t -channel subtraction	60	\pm	6	\pm	5				
$Z \rightarrow \ell^+ \ell^-$ (raw)	323	\pm	21	\pm	6	380	\pm	20	\pm 1
cross-section acceptance (%)	73.4				85.5				89.5
selection efficiency (%)	94.5				96.5				91.6
trigger efficiency (%)	100.0	—	0.2				98.3	\pm	0.6
$Z \rightarrow \ell^+ \ell^-$ (corrected)	466	\pm	30	\pm 13	469	\pm	25	\pm 5	653
$Z \rightarrow$ hadrons (corrected)	10385	\pm	103				9769	\pm	99
P_ℓ (%)	$4.48 \pm 0.30 \pm 0.12$			$4.80 \pm 0.26 \pm 0.05$			$4.70 \pm 0.21 \pm 0.11$		

the ratios of the numbers of lepton pairs to hadronic events where the latter were selected using the reconstructed charged tracks as described in section 3.

Table 4 shows these ratios (P) together with the branching ratios (B) and partial widths (Γ) deduced from the measured values of σ_{had}^0 and Γ_Z derived above, correcting for the effect of single photon exchange. Since the measured leptonic branching ratios are equal within errors, compatible with lepton universality, the average values $\Gamma_{\ell^+ \ell^-}$, $P_{\ell^+ \ell^-}$ and $B_{\ell^+ \ell^-}$ are also given. From these numbers the hadronic and invisible branching ratios are deduced following the procedure described in ref. [19]. They all agree very well with the standard model prediction.

The value of α_s can be checked by using Γ_{had} ; we find $\alpha_s = 0.22 \pm 0.10$, assuming the first order expansion $\Gamma_{\text{had}} = \Gamma_{\text{had}}^0 (1 + \alpha_s/\pi)$.

Since the branching ratios add to unity, the ratio of

Table 4
Measured ratios (P) of leptonic to hadronic Z decays, branching ratios (B) and partial widths (Γ) for the different decay modes.

Decay	P	B	Γ (MeV)
$Z \rightarrow e^+ e^-$	0.0448 ± 0.0031	0.0323 ± 0.0012	82.1 ± 3.4
$Z \rightarrow \mu^+ \mu^-$	0.0480 ± 0.0026	0.0346 ± 0.0023	87.9 ± 6.0
$Z \rightarrow \tau^+ \tau^-$	0.0470 ± 0.0024	0.0339 ± 0.0021	86.1 ± 5.6
$Z \rightarrow \ell^+ \ell^-$	0.0468 ± 0.0015	0.0330 ± 0.0006	83.9 ± 2.2
$Z \rightarrow$ hadrons		0.710 ± 0.015	1804 ± 44
$Z \rightarrow$ invisible		0.191 ± 0.014	495 ± 41

the invisible to the leptonic width is

$$\frac{\Gamma_{\text{inv}}}{\Gamma_{\ell^+ \ell^-}} = \frac{1}{B_{\ell^+ \ell^-}} - \frac{1}{P_{\ell^+ \ell^-}} - 3 = 5.78 \pm 0.39 ,$$

where $B_{\ell^+ \ell^-}$ and $P_{\ell^+ \ell^-}$ are the averaged leptonic branching ratios and the ratio of the leptonic to hadronic events, respectively, and Γ_{inv} and $\Gamma_{\ell^+ \ell^-}$ are the invisible and leptonic partial widths. Taking the value $\Gamma_{\text{inv}}/\Gamma_{\ell^+ \ell^-} = 1.99 \pm 0.01$, expected from the standard electroweak model, the number of neutrino types is 2.90 ± 0.19 . The importance of this result compared to the previous one, to which it is very strongly correlated, is that the ratio $\Gamma_{\text{inv}}/\Gamma_{\ell^+ \ell^-}$ is the only assumption from the standard model used in the analysis. The result is still valid if additional unexpected states yielding hadrons are present in Z decays.

The average leptonic partial width can be related to the effective weak mixing angle $\sin^2 \theta_w(M_Z)$ ⁴² in the following manner:

⁴² The effective weak mixing angle is defined as the ratio of the running QED and weak coupling constants evaluated at the Z mass:

$$\sin^2 \theta_w(M_Z) = \frac{e^2(M_Z)}{g^2(M_Z)} \approx 1 - \frac{M_W^2}{\rho M_Z^2} .$$

It is defined in different ways in the literature: $\sin^2 \theta_w(M_Z)_{\text{MS}}$ [21], $\sin^2 \theta_w^*(M_Z)$ [22], $\sin^2 \theta_w$ [23]. The numerical differences are insignificant.

$$\Gamma_{e^+e^-} = (1 + \kappa)$$

$$\times \frac{\alpha(M_Z) M_Z \{1 + [1 - 4 \sin^2 \theta_w(M_Z)]^2\}}{48 \sin^2 \theta_w(M_Z) \cos^2 \theta_w(M_Z)},$$

where $\kappa = (0.2 \pm 0.3)\%$ represents additional electro-weak effects [24] and $\alpha(M_Z)$ is the effective QED coupling constant.

From our measurement, $\Gamma_{e^+e^-} = 83.9 \pm 2.2$ MeV, we find

$$\sin^2 \theta_w(M_Z) = 0.231 \pm 0.008.$$

This determination is insensitive to assumptions on the top quark mass, the Higgs boson mass or the Higgs structure of the theory. It constitutes the most precise direct determination at this time of $\sin^2 \theta_w$ from neutral current couplings of leptons.

Acknowledgement

We would like to thank our colleagues of the LEP division for their outstanding performance in bringing the LEP machine into operation. Thanks also due to the many engineering and technical personnel at CERN and at the home institutes for their contribution towards ALEPH's success. Those of us from non-member states wish to thank CERN for its hospitality.

References

- [1] G.S. Abrams et al., *Phys. Rev. Lett.* 63 (1989) 724, 2173.
- [2] L3 Collab., B. Adeva et al., *Phys. Lett. B* 231 (1989) 509.
- [3] OPAL Collab., M.Z. Akrawy et al., *Phys. Lett. B* 231 (1989) 530.
- [4] DELPHI Collab., P. Aarnio et al., *Phys. Lett. B* 231 (1989) 539.
- [5] ALEPH Collab., D. Decamp et al., *Phys. Lett. B* 231 (1989) 519.
- [6] ALEPH Collab., ALEPH - A detector for electron-positron annihilation at LEP, *Nucl. Instrum. Methods*, to be published.
- [7] ALEPH Collab., D. Decamp et al., CERN preprint CERN-EP/89-139, submitted to *Phys. Lett.*
- [8] M. Bengtsson and T. Sjöstrand, *Phys. Lett. B* 185 (1987) 435.
- [9] F.A. Berends and R. Kleiss, *Nucl. Phys. B* 228 (1983) 737; M. Böhm, A. Denner and W. Hollik, *Nucl. Phys. B* 304 (1988) 687; F.A. Berends, R. Kleiss and W. Hollik, *Nucl. Phys. B* 304 (1988) 712.
- [10] H. Burkhardt, F. Jegerlehner, G. Penso and C. Verzegnassi, *Z. Phys. C* 43 (1989) 497.
- [11] E.A. Kuraev and V.S. Fadin, *Sov. J. Nucl. Phys.* 41 (1985) 466.
- [12] G. Altarelli and G. Martinelli, in: *Physics at LEP*, eds. J. Ellis and R. Peccei, CERN report CERN 86-02 (1986), Vol. I, p. 47.
- [13] F.A. Berends, G. Burgers and W.L. van Neerven, *Phys. Lett. B* 185 (1987) 395.
- [14] O. Niclosini and L. Trentadue, *Phys. Lett. B* 196 (1987) 551.
- [15] G. Burgers, in: *Polarization at LEP*, eds. G. Alexander et al., CERN report CERN 86-02 (1986), Vol I, p. 121.
- [16] A. Borelli, M. Consoli, L. Maiani and R. Sisto, CERN preprint CERN TH 5441 (1989).
- [17] A. Hofmann, private communication.
- [18] F.A. Berends et al., in: *Z Physics at LEP I*, eds. G. Altarelli, R. Kleiss and C. Verzegnassi, CERN report CERN 89-08 (1989), Vol. I, p. 89.
- [19] ALEPH Collab., D. Decamp et al., *Phys. Lett. B* 234 (1990) 399.
- [20] OPAL Collab., M.Z. Akrawy et al., *Phys. Lett. B* 235 (1990) 379.
- [21] W.J. Marciano and A. Sirlin, *Phys. Rev. Lett.* 46 (1981) 163.
- [22] D.C. Kennedy and B.W. Lynn, SLAC report SLAC Pub 4039 (1986), *Nucl. Phys. B* 322 (1989) 1.
- [23] M. Consoli, W. Hollik and F. Jegerlehner, in: *Z Physics at LEP I*, eds. G. Altarelli, R. Kleiss and C. Verzegnassi, CERN report CERN 89-08 (1989), Vol. I, p. 7.
- [24] D.C. Kennedy, B.W. Lynn, J.C. Im and R.G. Stuart, *Nucl. Phys. B* 321 (1989) 83.